

Erratum



2023; 13(15): 5597-5598. doi: 10.7150/thno.92042

Dual inhibition of PFKFB3 and VEGF normalizes tumor vasculature, reduces lactate production, and improves chemotherapy in glioblastoma: insights from protein expression profiling and MRI: Erratum

Junfeng Zhang^{1*}, Wei Xue^{1*}, Kai Xu¹, Liang Yi², Yu Guo¹, Tian Xie¹, Haipeng Tong¹, Bo Zhou¹, Shunan Wang¹, Qing Li³, Heng Liu⁴, Xiao Chen⁵, Jingqin Fang¹ and Weiguo Zhang^{1,6}

- 1. Department of Radiology, Daping Hospital, Army Medical University, Chongqing, 400042, China.
- 2. Department of Neurosurgery, Daping Hospital, Army Medical University, Chongqing, 400042, China.
- 3. Department of Oncology, Daping Hospital, Army Medical University, Chongqing, 400042, China.
- 4. Department of Radiology, PLA Rocket Force Characteristic Medical Center, Beijing, 100088, China
- 5. Department of Nuclear Medicine, Daping Hospital, Army Medical University, Chongqing, 400042, China.
- 6. Chongqing Clinical Research Center of Imaging and Nuclear Medicine, Chongqing, 400042, China.

*These authors contributed equally to this work.

© The author(s). This is an open access article distributed under the terms of the Creative Commons Attribution License (https://creativecommons.org/licenses/by/4.0/). See http://ivyspring.com/terms for full terms and conditions.

Published: 2023.11.17

Corrected article: Theranostics 2020; 10(16): 7245-7259. doi: 10.7150/thno.44427.

The authors regret that some inappropriate images were contained in Figure 3 from our paper. The immunostaining images of CD31/ α SMA, PIMO, collagen IV and LDHA at day 0, day 2, day 14 and day 25 from BEV+3PO, BEV and control groups were mis-pasted when choosing representative images from the large image data for typesetting. The correct version of Figure 3 appears below and these corrections made in this erratum do not affect the original data and conclusions. The authors apologize for any inconvenience that the errors may have caused.



Figure 3. Histological analysis of tumors at different time points after treatments. Representative images of double CD31 and α SMA-stained (A), collagenIV-stained (B), PIMO-stained (C), and LDHA-stained (D) GBM sections at different time points from different treatment groups (n = 120). The longitudinal assessments of MVD (E), pericyte coverage (F), collagenIV expression (G), tumor hypoxia (H), and LDHA expression (I) in the control, 3PO, BEV, and BEV+3PO groups at different time points. Scale bars, 50 µm (A, B), 100 µm (C, D). All data are means \pm SD. *P < 0.05, compared with control. *P < 0.05, compared with the BEV group.