

Myocardium-targeted transplantation of PHD2 shRNA-modified bone mesenchymal stem cells through ultrasound-targeted microbubble destruction protects the heart from acute myocardial infarction

Zhenxing Sun ^{1,2*}, Yuji Xie ^{1,2*}, Robert J. Lee ³, Yihan Chen ^{1,2}, Qiaofeng Jin^{1,2},
Qing Lv ^{1,2}, Jing Wang ^{1,2}, Yali Yang ^{1,2}, Yuman, Li ^{1,2}, Yu Cai ^{1,2}, Rui Wang ^{1,2},
Zhengyang Han ^{1,2}, Li Zhang ^{1,2}, Mingxing Xie ^{1,2}

¹Department of Ultrasound, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Wuhan 430022, China.

²Hubei Province Key Laboratory of Molecular Imaging, Wuhan 430022, China

³College of Pharmacy, The Ohio State University, Columbus, OH 43210, USA

Category of the Manuscript: Original Article

Total Word Count of the Manuscript:8029

*These authors contributed equally to this article.

 **Co-Corresponding author:**

1. Mingxing Xie, MD, Ph.D.; 1277 Jiefang Avenue, Wuhan, China. Tel: 86-27-85726430; Fax: 86-27-85726386. E-mail address: xiemx@hust.edu.cn.
2. Li Zhang, MD, Ph.D.; 1277 Jiefang Avenue, Wuhan, China. Tel: 86-27-85726430; Fax: 86-27-85726386. E-mail address: zli429@hust.edu.cn.




Figure S1. Characterization of BMSCs by microscopy and by flow cytometry.

A. The morphology observation of BMSCs under the optical microscopy. Bar, 50μm; B. Flow cytometry of BMSCs for mesenchymal CD29, 90, 34, and 45 markers. Unstained cells acted as control.