

Supporting information for

PET Imaging Facilitates Antibody Screening for Synergistic Radioimmunotherapy with a ¹⁷⁷Lu-labeled α PD-L1 Antibody

Jingyun Ren¹, Mengxin Xu², Junyi Chen², Jie Ding¹, Peipei Wang¹, Li Huo^{1*}, Fang Li^{1*}, Zhibo Liu^{2,3*}

¹Department of Nuclear Medicine, Peking Union Medical College Hospital, Chinese Academy of Medical Science & Peking Union Medical College, Beijing Key Laboratory of Molecular Targeted Diagnosis and Therapy in Nuclear Medicine, Beijing, China.

²Radiochemistry and Radiation Chemistry Key Laboratory of Fundamental Science, Beijing National Laboratory for Molecular Sciences, College of Chemistry and Molecular Engineering, Peking University, Beijing, 100871, China

³Peking University-Tsinghua University Center for Life Sciences, Beijing, 100871, China




Figure S1: Synthesis and radiolabeling of ^{89}Zr -DFO- α PD-L1.




Figure S2: Biodistribution of ^{89}Zr -DFO-Y001 (A) and ^{89}Zr -DFO-Y002 (B) in MC38 tumor bearing mice at 24 and 96 h post injection (n= 4).




Figure S3: Stability experiment of ^{177}Lu -DOTA-Y003 after incubation in PBS for 168 h.




Figure S4: Immunofluorescence for PD-L1 expression in 4T1 and MC38 cell lines.




Figure S5. Representative H&E staining of major organ tissues from mice with the indicated treatments (n = 3/group).