

Table S1. Primer sequences used in this study.

Gene	Forward Sequence (5'-3')	Reverse Sequence (5'-3')
<i>WNT1</i>	TGATACGCCAAAATCCGGGG	AGCCTCGGTTGACGATCTTG
<i>WNT2</i>	ACTCTCAGGACATGCTGGCT	ACGAGGTCATTTTTTCGTTGG
<i>WNT3</i>	TGTGAGGTGAAGACCTGCTG	AAAGTTGGGGGAGTTCTCGT
<i>LRP6</i>	CCCATGCCCCTGGTTCTACT	CCAAGCCACAGGGATACAGT
<i>MYC</i>	GAGCCTGCCTCTTTTCCACA	CAGCGACTCTGAGGAGGAAC
<i>BCL2</i>	CTGCACCTGACGCCCTTCACC	CACATGACCCACCCGAACTCAA AGA
<i>CCND1</i>	AAGTGCGTGCAGAAGGAGAT	TTAGAGGCCACGAACATGC
<i>BIRC5</i>	AGAACTGGCCCTTCTTGGAGG	CTTTTTATGTTCCCTCTATGGGGT C
<i>AXIN2</i>	ACTGCCACACGATAAGGAG	CTGGCTATGTCTTTGGACCA
<i>DNMT1</i>	CCATCAGGCATTCTACCA	CGTTCTCCTTGTCTTCTCT
<i>CD44</i>	CAATAGCACCTTGCCACAAT	AATCACCACGTGCCCTTCTATG G
<i>YAP1</i>	GCAACTCCAACCAGCAGCAA	CGCAGCCTCTCCTTCTCCAT
<i>SNAI2</i>	TGTTGCAGTGAGGGCAAGAA	GACCCTGGTTGCTTCAAGGA
<i>TWIST1</i>	GGACAAGCTGAGCAAGAT	CTCTGGAGGACCTGGTAG
<i>VIM</i>	CGGCTGCGAGAGAAATTGC	CCACTTTCCGTTCAAGGTCAAG
<i>CDH2</i>	GGTGGAGGAGAAGAAGACCAG	GGCATCAGGCTCCACAGT
<i>NANOG</i>	CCTCCTCCATGGATCTGCTTATT CA	CAGGTCTTACCTGTTTGTAG
<i>POU5F1</i>	TGCAGCAGATCAGCCACATC	CTCGGACCACATCCTTCTCG
<i>SOX2</i>	AACCAGCGCATGGACAGTTA	ATCATGCTGTAGCTGCCGTT
<i>ACTB</i>	TCATTCCAAATATGAGATGCGTT G	TAGAGAGAAGTGGGGTGGCT
<i>RN18S</i>	CGCCGCTAGAGGTGAAATTC	TTGGCAAATGCTTTCGCTC




Figure S1. Regulation of mRNA expression of Wnt ligands in 17-AAG-resistant cells. Real-time PCR analysis showing changes in mRNA expression of Wnt ligands in 17-AAG-resistant (17AAG^R) H1299 and A549 cells by comparison with those in the corresponding parental (P) cells. The bars represent mean \pm SD; * $P < 0.05$, ** $P < 0.01$, and *** $P < 0.001$, as determined by a two-tailed Student's *t*-test in comparison with the indicated control.